



**CULLMAN**  
REGIONAL

## PHYSICIAN'S ORDERS

NAME:  
ROOM NO:  
(ADDRESS)  
HOSP. NO.  
PHYSICIAN

Another brand of drug identical in form and content may be dispensed unless checked. ☐

### OB/GYN – Cullman Anesthesiology Order Set

#### ☐ Epidural Anesthesia

1. Hydrate with 1000 cc IV LR before epidural.
2. If IV infusing more than 2 hours, hydrate with 500 cc LR bolus.
3. After test dose or boost, check BP & Pulse q 2 minutes x 4, q 5 minutes x 4, then q 15 minutes.
4. Check level of anesthesia (use alcohol pad and check for coolness q 30 minutes)  
T10 = Umbilicus  
T6 = Xiphoid  
T4 = Nipple
5. If there is a rapid fall in BP more than 30 mmHg or a fall to 90 mmHg systolic, then RN may give Ephedrine 5 mg. IV Stat and call MDA.
6. If sensory level is consistently above T10 during first stage of labor, notify MD Anesthesiologist.
7. If patient is unable to push, Notify MD Anesthesiologist
8. Emergency/Unscheduled C-Section
  - a. Pepcid 20 mg IV
  - b. Bicitra 30 cc Po
9. Naropin 0.2% via pump per anesthesia
10. Fentanyl 100 Mcg administered by anesthesia

MD Signature: \_\_\_\_\_ Date & Time: \_\_\_\_\_

**Please use Ball Point Pen ONLY**

**DO NOT USE: U IU QD QOD MS MSO4 MgSO4**