



**CULLMAN
REGIONAL**

PHYSICIAN'S ORDERS

NAME:
ROOM NO:
(ADDRESS)
HOSP. NO.
PHYSICIAN

Date & Time	Another brand of drug identical in form and content may be dispensed unless checked. <input type="checkbox"/>
	Micheal J. Lunsford, M.D. Adult patient (>18 years old) Standing Orders
1. Notify Dr. Lunsford at 0700 of admission if admitted overnight	
2. Vital signs per unit routine	
3. Notify MD for temp > 101.5, HR > 150 or < 50, Systolic BP > 220 or diastolic BP > 100 or < 40, O2 Sat < 90%, or other acute changes	
4. Diet: 2000 Cal ADA diet if diabetic, cardiac prudent diet if patient with coronary disease, renal diet if patient with renal disease	
5. Tylenol 650 mg Po/PR every 4 hr PRN mild pain (scale 1-3) or temp > 101.5	
6. Zofran 4 mg IV every 4 hr PRN nausea	
7. Phenergan 25 mg PR every 4 hr PRN nausea if no IV access	
8. Lactulose 30 ml Po every 6 hr PRN constipation	
9. If no results from Lactulose, then give 1 Dulcolax supp PR x 1	
10. Maalox 30 ml Po every 6 hr PRN indigestion	
11. Imodium 2 tablets initially Po for severe diarrhea, then 1 Po every 4 hr PRN diarrhea	
12. Robitussin DM 10 ml Po every 4 hr PRN cough	
13. Trazodone 25 mg Po every hs PRN for insomnia	
14. May use 1% Lidocaine 2 ml with 1 gm Rocephin or 1% Lidocaine 3 ml with 2 gm Rocephin for IM injections.	
15. Obtain EKG and give 0.4 mg Nitroglycerin SL PRN x 1 for new angina, notify MD if pain not relieved promptly. Hold Nitroglycerin for Systolic BP < 90 mmHg	
16. May insert Foley for signs of urinary retention; implement Foley Catheter Removal Protocol	
17. Accuchecks q 0700, 1100, 1600, 2100 on all diabetics	
18. Automatically pull old charts on patients admitted	
19. T pump PRN	
20. Clonidine 0.1 mg Po PRN x 1 for Systolic BP > 200 (except if patient has new CVA this admit). May repeat in 1 hour if Systolic BP > 200 x 1, then notify MD if Systolic BP remains > 180.	
21. If current order for Albuterol treatment, may give one extra Albuterol 2.5 mg nebulized treatment for wheezing/short of breath (during 11 pm-7 am). May check saturation per respiratory and notify MD if SaO2 < 88% or no results from treatment. While calling MD, patient may be placed on O2 NC @ 2 L/m for saturation less than 88%.	
22. Notify MD for K+ < 3.1.	
MD Signature: _____ Date & Time: _____	

Cullman Regional

Please use Ball Point Pen ONLY

Physician's Orders

DO NOT USE: U IU QD QOD MS MSO4 MgSO4